





































42.16.4.2.8.4	Method 4:	Synthesis from Methyl Bis(4-nitrophenyl) Phosphite with Subsequent Sulfurization or Selenization	480
42.16.4.2.8.5	Method 5:	Lithium Iodide/ <i>tert</i> -Butyl Hydroperoxide Mediated Oxidative Cross Coupling of <i>O,O</i> -Dimethyl Phosphonothioate with Phenols	482
42.16.4.2.9		<i>O,O,S</i> -Trialkyl/ <i>O,O,S</i> -Triaryl Phosphorodithioates	483
42.16.4.2.9.1	Method 1:	Sulfurization of Trivalent Phosphorus Compounds	483
42.16.4.2.9.2	Method 2:	Alkylation of Phosphorodithioates	485
42.16.4.2.9.2.1	Variation 1:	Selective Ring Opening of Vinyloxetanes with <i>O,O</i> -Dialkyl <i>S</i> -Hydrogen Phosphorodithioates	485
42.16.4.2.9.2.2	Variation 2:	Reaction with Nonactivated Alkenes in the Presence of Zinc(II) Chloride	486
42.16.4.2.9.2.3	Variation 3:	Solvent-Free Approach Using Alumina	487
42.16.4.2.9.2.4	Variation 4:	From Monoterpene Alcohols via <i>S</i> -Alkylation or <i>S</i> -Acylation	488
42.16.4.2.9.3	Method 3:	Alcoholysis of Phosphorodichloridodithioates	489
42.16.4.2.9.4	Method 4:	Chemoselective <i>S</i> -Alkylation of <i>O,O</i> -Dialkyl Phosphorodithioates	490
	<b>Author Index</b>		499
	<b>Abbreviations</b>		523